

# Oral Candidal and Streptococcal carriage in Down syndrome patients

Gouse Mohiddin,  
Aravindha Babu  
Narayanaswamy<sup>1</sup>,  
Masthan K. M. K.<sup>1</sup>,  
Anitha Nagarajan<sup>1</sup>,  
Abikshyeet Panda,  
Shyam Sundar Behura

Department of Oral and Maxillofacial Pathology, Kalinga Institute of Dental Sciences, Bhubaneswar, Odisha, <sup>1</sup>Department of Oral and Maxillofacial Pathology, Sree Balaji Dental College and Hospitals, Chennai, Tamil Nadu, India

**Address for correspondence:**

Dr. N. Anitha, Department of Oral and Maxillofacial Pathology, Sree Balaji Dental College and Hospitals, Chennai - 600 010, Tamil Nadu, India. E-mail: [dranitha.nnr@gmail.com](mailto:dranitha.nnr@gmail.com)

## Abstract

**Aim:** We aimed to evaluate the prevalence of *Candida* and Streptococci species in the oral cavity of Down syndrome patients. **Materials and Methods:** 50 children/adolescents with Down syndrome with a karyotype of 47 XX, 21+ (female) and 47 XY, 21+ (male), and 50 normal children/adolescents were included in our study. Oral swab/saliva was used to culture and identify *Candida* and Streptococci species based on gram and periodic acid schiff staining. **Results:** Of the 50 study group samples, which were cultured, 37 (74%) showed growth of *Candida* colonies, whereas in the 50 control samples only 18 (36%) were positive for *Candida* growth. In 4 Sabouraud's dextrose agar culture slopes of the study group, more than one morphological type of colonies were observed. 23 out of 50 samples in our study group had *Streptococcus viridans* colonies. In the 23 samples positive for Streptococci 16 had many streptococcal colonies, and 7 had few streptococcal colonies in the primary culture. 32 out of 50 samples from the control group had *S. viridans* colonies. In these 32 samples positive for Streptococci, 29 had predominantly streptococcal colonies while 3 had few streptococcal colonies in the primary culture. **Conclusion:** The oral cavity is an environment heavily colonized by microorganisms, however, the Down syndrome patients run a greater risk of having opportunistic infections especially from *Candida* species. Hence to improve the quality of life of an individual with Down syndrome, it is necessary to diagnose and treat these infections by more frequent oral microbial assessment.

**Key words:** *Candida*, culture, Down syndrome, microbes, *Streptococcus*

## INTRODUCTION

Down syndrome (trisomy 21) is the most common and best known of all malformation syndromes. It is an easily recognized congenital, autosomal anomaly characterized by generalized growth deficiency and mental deficiency affecting 1 in 600-1 in 1000 live births. It accounts for about one-third of all mentally handicapped children.<sup>[1]</sup> Many systemic diseases are responsible for mortality of infants and children with Down syndrome. A compromised immune system is characteristic of these individuals, and it

is this immunodeficiency that contributes to an increased susceptibility to infections. Various microorganisms are generally seen as normal commensals in the oral cavity, and these may become pathogenic under favorable conditions. *Candida* is one such endogenous pathogen, the prevalence of which has been reported to be 45-65% in healthy children<sup>[2]</sup> and 30-45% in healthy adults.<sup>[3]</sup> Opportunistic infections occur mainly when the host defenses are inadequate, candidiasis is the most common among them. Streptococci form a large portion of the resident oral micro flora. In the general population, half the isolates from the tongue and saliva are Streptococci, but in Down syndrome individual's oral streptococcal levels are relatively lower.<sup>[4]</sup> This probably contributes to a disturbance in the normal microbial homeostasis and promotes the growth of other commensal organisms like *Candida*. Here, we evaluated the candidial/Streptococci carriage and the various species of *Candida*/Streptococci isolated from the Down syndrome patients and correlated it with normal oral micro flora from health subjects.

| Access this article online  |  |
|---|--|
| Quick Response Code:<br> | Website:<br><a href="http://www.jnsbm.org">www.jnsbm.org</a> |
|   | DOI:<br>10.4103/0976-9668.159983                             |

## MATERIALS AND METHODS

Fifty Down syndrome individuals were selected for this study based on a karyotype of 47 XX, 21+ (female) and 47 XY, 21+ (male). A total of 50 normal children/adolescents were selected as a control group. To isolate *Candida* and Streptococci, swab/saliva was used to culture and smear preparation for Gram stain and periodic acid schiff (PAS) stain.

### Culture for *Candida*

Samples were taken from the oral cavity (dorsum of tongue) for each individual using a sterile cotton wool swab. Sabouraud's dextrose agar (SDA) slope with antibiotic (gentamicin/actidione) was used for the isolation of *Candida*. Antibiotics were used in order to eliminate bacterial contamination. If colonies were white or cream to yellow, dull/shiny, pasty yeast like a smear was prepared from the colony, with a drop of saline. Smear was then air-dried, heat fixed, and stained with crystal violet (simple stain) to confirm the presence of yeast. The isolates were then subjected to the following tests for identification of various species of *Candida*:

1. Germ tube test
2. Morphology on corn meal agar — for chlamydospore formation.
3. Carbohydrate fermentation test
4. Carbohydrate assimilation test
5. CHROMagar *Candida* culture characteristics

### Smear and Gram stain

Smears taken from the oral cavity (tongue) were air-dried, heat fixed, and stained.

## RESULTS

The study included a total of 100 (50 Down syndrome, 50 controls) subjects of both sexes. The study group of 50 Down syndrome individuals comprised of 37 males and 13 females in the age range of 3-22 years and the mean age of 11.6 years. The control group of 50 normal individuals comprised of 34 males and 16 females in the age range of 3-18 years and a mean age of 9.5 years [Tables 1 and 2].

Of the 50 study group samples, which were cultured, 37 (74%) showed growth of *Candida* colonies, whereas in the 50 control samples only 18 (36%) were positive for *Candida* growth. In 4 SDA culture slopes of the study group, more than one morphological type of colonies were observed [Figure 1], and the colonies were considered as different isolates and subjected to further testing. A total of 42 isolates (1 isolate in 33 samples, 2 isolates in 3 samples, and 3 isolates in 1 sample) were obtained from the 37 culture positive samples, in the study group and 18 isolates were

obtained from the 18 culture positive samples in the control group (1 isolate each in 18 samples) [Table 3].

Smears taken from the oral cavity (tongue), for both the study group (50) and the control group (50) were stained with Gram stain [Figure 2] and PAS stain [Figure 3]. A tenfold increase in the positive staining for *Candida* was observed in the study group when compared to the control group [Table 4].

Of the 42 isolates from the 37 culture positive samples, 31 were germ tube positive, whereas all the 18 culture positive samples of the control group were positive for germ tube formation [Figure 4]. Hence, 31 of the 42 isolates were *Candida albicans* in the study group, and all 18 isolates of the



**Figure 1:** Sabouraud's dextrose agar slope showing more than one type of colony (white and cream)

**Table 1: Distribution of patients according to sex**

| Sex    | Study group | Control group |
|--------|-------------|---------------|
| Male   | 37          | 34            |
| Female | 13          | 16            |
| Total  | 50          | 50            |

**Table 2: Distribution of patients according to age**

| Age               | Study group | Control group |
|-------------------|-------------|---------------|
| Age range (years) | 3-22        | 3-18          |
| Mean age (years)  | 11.6        | 9.5           |

**Table 3: Candidal growth pattern in the study group and control group**

| Growth pattern                     | Study group | Control group |
|------------------------------------|-------------|---------------|
| Total number of samples            | 50          | 50            |
| Candidal growth                    | 37          | 18            |
| Total number of isolates           | 42          | 18            |
| Samples with more than one isolate | 4           | 0             |

**Table 4: Distribution of positive candidal staining in study group and control group**

| Group   | Number (%)       |               |
|---------|------------------|---------------|
|         | Gram stain steps | PAS stain car |
| Study   | 10 (20)          | 12 (24)       |
| Control | 1 (2)            | 1 (2)         |

PAS: Periodic acid schiff

control group were *C. albicans*. This was further confirmed by chlamydospore formation on corn meal agar [Figure 5]. All the isolates that were germ tube positive produced chlamydospores on corn meal agar further confirming *C. albicans*. Of the 42 isolates and 37 culture positive samples, 31 isolates were *C. albicans*, 6 were *Candida Tropicalis*, and 5 were either *Candida parapsilosis*/*Candida krusei*/*Candida glabrata*. Of the 18 control samples, which were culture positive, all 18 were *C. albicans*. Further confirmation and speciation was done by the carbohydrate assimilation test and supplemented with growth on CHROMagar *Candida*.

### Carbohydrate assimilation test

This test was done with sugars: Glucose, maltose, sucrose, lactose, galactose, cellulose, and trehalose [Table 5].

Of the 37 samples that were culture positive in the study group, 31 (84%) samples had isolates of *C. albicans*, 6 (16%)

samples had isolates of *C. tropicalis* and 5 (14%) had isolates of *C. parapsilosis*. 4 samples of the study group were colonized with more than one type of species. Of the 18 control samples that were culture positive all 18 isolates were *C. albicans*.

### Chromagar Candida

Growth of all culture positive samples on CHROMagar *Candida* was observed for the characteristic colony color [Table 6].

In the CHROMagar *Candida*, study sample showed 31 isolates of *C. albicans*, 6 of *C. tropicalis*, and 5 of *C. parapsilosis* [Figure 6]. In the control samples, all 18 of the isolates were *C. albicans*, thus confirming the earlier results.

### Culture for identification of streptococci

Saliva samples collected from both the study group (50%) and the control group (50) after inoculation on chocolate




Figure 2: Gram-stained smear showing yeasts and pseudohyphae



Figure 3: Periodic acid schiff stained smear showing candidal hyphae and yeasts




Figure 4: Germ tube formation by *Candida albicans*




Figure 5: Chlamydospore formation by *Candida albicans*

Table 5: Carbohydrate assimilation reactions for different *Candida* species

| Candida species  | Study group: 37 culture positive samples (42 isolates) |         |         |         |           |            |           | Number of isolates |
|--|--|---------|---------|---------|-----------|------------|-----------|--------------------|
|  | Glucose  | Maltose | Sucrose | Lactose | Galactose | Cellobiose | Trehalose |                    |
| <i>C. albicans</i>                                       | +  | +       | +       | -       | +         | -          | +         | 31                 |
| <i>C. tropicalis</i>                                     | +  | +       | +       | -       | +         | +          | +         | 6                  |
| <i>C. parapsilosis</i>                                   | +  | +       | +       | -       | +         | -          | +         | 5                  |
| Control group: 18 culture positive samples (18 isolates) |  |         |         |         |           |            |           |                    |
| <i>C. albicans</i>                                       | +  | +       | +       | -       | +         | -          | +         | 18                 |

*C. albicans*: *Candida albicans*, *C. tropicalis*: *Candida tropicalis*, *C. parapsilosis*: *Candida parapsilosis*



agar, and incubation at 37°C for 24 h, the growth of alpha-hemolytic colonies were identified [Figure 7]. The colonies were checked for optochin resistance to classify it as *Streptococcus viridans* group [Table 7].

The growth of alpha hemolytic *S. viridans* colonies in the primary culture was classified as in Table 8.

Twenty-three out of 50 samples from study group had *S. viridans* colonies. In the 23 samples positive for Streptococci 16 had many streptococcal colonies, and 7 had few streptococcal colonies in the primary culture. Of the 50 control samples, 32 had *S. viridans* colonies. In the 32 samples, which were positive for Streptococci, 29 had predominantly streptococcal colonies while 3 had few streptococcal colonies in the primary culture.

### Association of *Candida* with streptococci

The presence of *Candida* with Streptococci was compared in both the study group and the control group [Table 9].

Candidal and streptococcal colonization was seen in 14 (28%) individuals in the study group and 6 (12%) individuals in the control group. *Candida* colonization alone in the absence of streptococcal colonization was seen in 23 (46%) individuals of the study group and 12 (24%) of the control group. Thus showing that candidal colonization occurred to a greater extent in the absence of Streptococci ( $P < 0.01$ ,  $\chi^2 = 2.96$ ).

Our results showed that in the study group (74%) *Candida* colonization was comparatively higher when compared, with the control group (36%). Of the 37 culture-positive samples, 42 isolates were obtained, four samples had more than one isolate (one sample with three isolates, three samples with two isolates). The species isolated in the study group out of the 37 culture-positive samples were predominantly *C. albicans* (84%) followed by *C. tropicalis* (16%) and *C. parapsilosis* (14%). In the control group, only *C. albicans* were isolated. Thus, out of the total 50 Down

syndrome subjects, 31 (62%) showed colonization with *C. albicans*, 6 (12%) with *C. tropicalis*, and 5 (10%) with *C. parapsilosis*. In the comparative study of *Candida* and Streptococci, *Candida* colonization was predominantly seen in the absence of Streptococci.

**Table 6: CHROMagar *Candida* culture characteristics for different *Candida* species**

| Study group: 37 culture positive samples (42 isolates)   |                        |                    |
|--|------------------------|--------------------|
| Colony color   | Species                | Number of isolates |
| Green  | <i>C. albicans</i>     | 31                 |
| Blue   | <i>C. tropicalis</i>   | 6                  |
| Cream  | <i>C. parapsilosis</i> | 5                  |
| Control group: 18 culture positive samples (18 isolates) |                        |                    |
| Green  | <i>C. albicans</i>     | 18                 |

*C. albicans*: *Candida albicans*, *C. tropicalis*: *Candida tropicalis*, *C. parapsilosis*: *Candida parapsilosis*

**Table 7: Growth pattern of *Streptococcus***

| Group   | Total cases | Alpha hemolytic colonies positive | Optochin resistance | Percentage |
|---------|-------------|-----------------------------------|---------------------|------------|
| Study   | 50          | 23                                | 23                  | 46         |
| Control | 50          | 32                                | 32                  | 64         |

**Table 8: Distribution of viridans group of Streptococci in control group and study group**


| Alpha hemolytic colonies | Study group | Control group |
|--------------------------|-------------|---------------|
| Many                     | 16          | 29            |
| Few                      | 7           | 3             |
| None                     | 27          | 18            |

**Table 9: Comparison of candidal and streptococcal growth patterns in study group and control group**

| Growth pattern                          | Number (%)       |                    |
|---|------------------|--------------------|
|   | Study group (50) | Control group (50) |
| Candidal and streptococcal colonization | 14 (28)          | 6 (12)             |
| <i>Candida</i> colonization alone       | 23 (46)          | 12 (24)            |



**Figure 6:** CHROMagar *Candida* showing green colonies of *Candida albicans* and cream colonies of *Candida parapsilosis*



**Figure 7:** Primary culture showing predominantly alpha hemolytic colonies

### Oral manifestations

The oral findings of all the children and adolescents of the study group were recorded [Figures 8-10]. Table 10 shows in an ascending order the frequency of various findings.

In the 50 Down syndrome individuals, 17 (34%) of them showed an erythematous or a pseudomembranous lesion on the tongue (12 - erythematous, 5 - pseudomembranous), indicating possible oral candidiasis.

### DISCUSSION

Down syndrome individuals are reported to exhibit an increased susceptibility to infections due to various factors such as increased activity of superoxide dismutase,<sup>[5,6]</sup> increased activity of glutathione peroxidase,<sup>[6]</sup> decreased levels of myeloperoxidase in leukocytes,<sup>[7]</sup> defective neutrophil chemotaxis,<sup>[8]</sup> abnormal serum immunoglobulin patterns<sup>[9]</sup> and disturbed cell-mediated and humoral immunity.<sup>[10]</sup>

The most common opportunistic infection affecting the oral cavity is candidiasis. The present study determined the prevalence of oral *Candida* carriage in Down syndrome children and adolescents. Of the 37 individuals in the study group who exhibited *Candida* colonization, only 14 showed colonization with Streptococci while 23 remained negative and showed no streptococcal colonization. Thus, we concluded that the majority of the individuals showed *Candida* colonization when there was no concomitant streptococcal colonization. The increased *Candida* carriage in Down syndrome individuals could probably be attributed to the relatively lower streptococcal levels. The increased susceptibility to *Candida* infection may be a result of the disturbance in the microbial homeostasis. However, other factors such as a compromised immune system, poor oral hygiene, malnourishment and the like cannot be completely



Figure 8: 12-year-old male showing fissured tongue




Figure 9: High arched palatal vault



Figure 10: Angular cheilitis with prognathic mandible

ruled out and are also expected to play a significant role. It may be also possible that the *Candida* and Streptococci may produce mutually inhibitive factors in the form of toxins or enzymes, thus producing this inverse relationship.

The most common predisposing factor resulting in candidiasis is the presence of a suppressed immune status, thus this infection is seen in a higher frequency in Down syndrome subjects, where a compromised immune status is the hallmark.<sup>[11]</sup> In addition to this, a variety of local factors may also play a role, the most common being an altered intra oral milieu. As observed in this study, it may be assumed that the presence of microbial imbalance in the oral cavity and the occurrence of fissured tongue could be some major local factors contributing to the increased *Candida* carriage in Down syndrome individuals.

### CONCLUSION

The oral cavity is an environment heavily colonized by microorganisms, which are generally in harmony with each other. This relationship seems to be related to the immune status of the individual. The Down syndrome subjects run a greater risk of having opportunistic infections with a possibility of systemic spread, and the associated morbidity

**Table 10: Distribution of oral findings in down syndrome subjects**

| Manifestation                     | Number of subjects affected | Percentage prevalence |
|-----------------------------------|-----------------------------|-----------------------|
| Open mouth with tongue protrusion | 47                          | 94                    |
| Angle's class III malocclusion    | 33                          | 66                    |
| Macroglossia                      | 28                          | 56                    |
| Large lingual papilla             | 24                          | 48                    |
| Fissured tongue                   | 19                          | 38                    |
| Angular cheilitis                 | 18                          | 36                    |
| Narrow high arched palate         | 13                          | 26                    |
| Diastema                          | 12                          | 24                    |
| Partial anodontia                 | 8                           | 16                    |
| Crowding                          | 8                           | 16                    |
| Peg-shaped teeth                  | 6                           | 12                    |
| Enamel hypoplasia                 | 4                           | 8                     |

and mortality is also higher. Hence, to improve the quality-of-life of an individual with Down syndrome, it becomes necessary to diagnose and treat these infections when they still remain localized.

## REFERENCES

1. Regezi JA, Sciubba JJ, Jordan RC. Oral Pathology: Clinical Correlations. 4<sup>th</sup> ed. USA: Elsevier Science; 2003. p. 358-9.
2. Berdicevsky I, Ben-Aryeh H, Sazargel R. Oral *Candida* in children. Oral Surg Oral Med Oral Pathol 1980;57:37-40.
3. Akpan A, Morgan R. Oral candidiasis. Postgrad Med J 2002;78: 455-9.
4. Lee SR, Kwon HK, Song KB, Choi YH. Dental caries and salivary immunoglobulin A in Down syndrome children. J Paediatr Child Health 2004;40:530-3.
5. Mattei JF, Baeteman MA, Baret A, Ardisson JP, Rebuffel P, Giraud F. Erythrocyte superoxide dismutase and redox enzymes in trisomy 21. Acta Paediatr Scand 1982;71:589-91.
6. Sinet PM, Michelson AM, Bazin A, Lejeune J, Jerome H. Increase in glutathione peroxidase activity in erythrocytes from trisomy 21 subjects. Biochem Biophys Res Commun 1975;67:910-5.
7. Björkstén B, Marklund S, Hägglöf B. Enzymes of leukocyte oxidative metabolism in Down's syndrome. Acta Paediatr Scand 1984;73: 97-101.
8. Izumi Y, Sugiyama S, Shinozuka O, Yamazaki T, Ohyama T, Ishikawa I. Defective neutrophil chemotaxis in Down's syndrome patients and its relationship to periodontal destruction. J Periodontol 1989;60: 238-42.
9. Avanzini MA, Söderström T, Wahl M, Plebani A, Burgio GR, Hanson LA. IgG subclass deficiency in patients with Down's syndrome and aberrant hepatitis B vaccine response. Scand J Immunol 1988;28: 465-70.
10. Annerén G, Magnusson CG, Lilja G, Nordvall SL. Abnormal serum IgG subclass pattern in children with Down's syndrome. Arch Dis Child 1992;67:628-31.
11. Scully C. Down's syndrome and dentistry. Dent Update 1976;3:193-6.

**How to cite this article:** Mohiddin G, Narayanaswamy AB, Masthan K, Nagarajan A, Panda A, Behura SS. Oral Candidal and Streptococcal carriage in Down syndrome patients. J Nat Sc Biol Med 2015;6:300-5.

**Source of Support:** Nil. **Conflict of Interest:** None declared.

